Dermatology Enrollment Form – Page 1 of 2

TwelveStone Health Partners Fax Referral To: (200) 223-4063



| Patient Name: | Patient Name: HEALTH PARTNERS ** | | | | | | | | | | |
|--|----------------------------------|--|---|--|---------|------------------------|--|--|--|--|--|
| | | | Direct Phone: (615 | 5) 2/8-3350 | | | | | | | |
| Date of Birth: | | | Toll Free: (844) 8 | 033-0012 | | | | | | | |
| PREVIOUS ADMINISTRATION | | | | | | | | | | | |
| • | ovide the following inf | | | If NO, please indicate desired location for first dose: | | | | | | | |
| | te: | | | Physician's Office | | | | | | | |
| Next Infusion Da | te: | | | TwelveStone Infusion Suite | | | | | | | |
| | | | L | ☐ TwelveStone Home Infusion☐ Other: | | | | | | | |
| | | | | Desired Start Date: | | | | | | | |
| DIAGNOSIS | | | | | | | | | | | |
| Description ICD-10Code | | | | | | | | | | | |
| | | | | | | | | | | | |
| OTHER REQUIRED DOCUMENTATION (Please attach documents as needed) | | | | | | | | | | | |
| | | | | | | | | | | | |
| | ☐ This s | signed order form | ☐ History and Phy | rsical TB and Hep B Doo linical progress notes, lab work (includi | | | | | | | |
| | Patient Demographics | and Insurance Inform | | unction tests and any other tests suppo | • | | | | | | |
| CLINICAL INFORMATION (Please attach all clinical information, lab results and other medical history documents) | | | | | | | | | | | |
| Patient Weigh | nt· Ka/ | Lbs Height: | Inches/CM RSA: | Allergies: | | | | | | | |
| r atient weigi | | | · | | . – | | | | | | |
| | Line Access: | ☐ PIV ☐ | PICC (SL DL TL) POR | T (Huber sizeGaugeLength |) 🗆 🤄 | Sub-Q | | | | | |
| | | CURREN | TLY RECEIVING AND/OR PF | RIOR FAILED THERAPIES: | | | | | | | |
| ☐ Biologics: ☐ | Cimzia □ Cosentyx □ | ☐ Enbrel ☐ Humira | ☐ Orencia ☐ Remicade ☐ R | ituxan □ Simponi □ Stelara | | | | | | | |
| ☐ Methotrexate | e □ Soriatane □ CYA | A 🗆 PUVA/UVB 🗆 | Topicals 🗆 Other | | | | | | | | |
| Length of Treatm | nent | | | | | | | | | | |
| | | | | | | | | | | | |
| Contradicted Med | dications: | | | | | | | | | | |
| Reason: | | | | | | | | | | | |
| | | | | | | | | | | | |
| MEDICATION | | | | | REFILLS | LAB & ANCILLARY ORDERS | | | | | |
| ☐ Cosentyx | ☐ 150mg/ml Pen | ☐ Initiation - Injec | t 300mg SQ at 0,1,2,3,4 and th | nen every 4 weeks thereafter | | | | | | | |
| □ Cosemyx | ☐ 150mg/ml PFS | ☐ Maintenance - I | nject 300mg SQ every 4 weeks | 5 | | | | | | | |
| ☐ Humira | ☐ 40mg Pen | ☐ Initiation - Inject | ☐ Initiation - Inject 80mg SQ followed by 40mg every other week starting 1 week after | | | | | | | | |
| | ☐ 80mg Pen | ☐ Maintenance - Ir | ☐ Maintenance - Inject 300mg SQ every 4 weeks | | | | | | | | |
| | | ☐ Initiation - 100m | ng SQ at 0, 4 weeks, and then eve | ery 8 weeks thereafter. | | | | | | | |
| ☐ Tremfya | 100mg/ml PFS | | nject 100mg SQ every 8 weeks | · | | | | | | | |
| | ☐ 45mg PFS | ☐ Initiation (< 100kg) – Inject 45mg SQ week 0, and 4, then 45mg every 12 weeks thereafter | | | | | | | | | |
| □ Stolara | | ☐ Maintenance (<100kg) - Inject 45mg SQ every 12 weeks | | | | | | | | | |
| ☐ Stelara | □ 90 mg PES | Initiation (> 100kg) Inject 00mg 50 week 0 and 4 then 00mg every 12 weeks | | - | | | | | | | |
| | □ 90 mg PFS | ☐ Initiation (> 100kg) – Inject 90mg SQ week 0, and 4, then 90mg every 12 weeks thereafter | | | | | | | | | |
| | | ☐ Maintenance (> | 100kg) - Inject 90mg every 12 w | reeks | | | | | | | |
| ☐ Dupixent | 300mg/2ml PFS | ☐ Initiation - Inject | 600mg SQ day 1, then 300mg o | on day 15, then 300mg every other week | | | | | | | |
| | | ☐ Maintenance - Ir | nject 300mg SQ every other wee | ek | | | | | | | |
| ☐ Odomzo | 200mg Capsule | Take 1 (one) capsule by mouth daily on an empty stomach | | | | | | | | | |
| ☐ Otrexup | | Inject mg SQ | weekly (10-25mg usual dose) | | | | | | | | |
| □ Otezla | ☐ Starter Pack | ☐ Initiation - Titra | te dose up to 30mg PO BID sta | orting with 10mg qAM | | | | | | | |
| | ☐ 30mg Tablets | ☐ Maintenance - Take 1 (one) tablet my mouth twice daily | | | | | | | | | |
| ☐ Cimzia | 200mg/ml PFS | | : 400mg SQ (2 injections) at wee | · | | | | | | | |
| | | | nject 200mg SQ every other wee | | | | | | | | |
| | | | enance - Inject 400mg SO (2 inje | | 1 | | | | | | |

Dermatology Enrollment Form - Page 2 of 2

| Date: | |
|----------------|--|
| Patient Name: | |
| Date of Birth: | |

TwelveStone Health Partners Fax Referral To: (800) 223-4063

Direct Phone: (615) 278-3350 Toll Free: (844) 893-0012



| MEDICATION | DOSE | DIRECTIONS | | REFILLS | LAB & ANCILLARY ORDERS | | | | |
|--|---------------------------|---|---|---------|-------------------------------|--|--|--|--|
| ☐ Siliq | | ☐ Initiation - 210mg SQ at 0,1,2 weeks, and then every 2 weeks thereafter | | | | | | | |
| | 210mg PFS | ☐ Maintenance - Inject 210mg SQ every 2 weeks | | | | | | | |
| ☐ Remicade | | ☐ Initiation - Inject mg IV infusion at (5mg/kg) | 0,2,6 weeks and then every 8 weeks thereafter | | | | | | |
| | | ☐ Maintenance - Infuse mg IV every 8 weeks (5mg/kg) | | | | | | | |
| ☐ Rasuvo | | Inject mg SQ weekly (7.5-30mg usual dose) | | | | | | | |
| □ Enbrel | ☐ Sureclick 50mg/ml | ☐ Initiation - Inject 50mg SQ twice weekly x 3 months; then 50mg weekly thereafter (adult dosing) | | | | | | | |
| | ☐ 25mg/ml PFS | ☐ Maintenance - Inject 50mg SQ weekly | | | | | | | |
| | ☐ 50mg/ml PFS | ☐ Maintenance - Inject 0.8mg/kg (mg) SQ once weekly (Pediatric dosing) | | | | | | | |
| ☐ Simponi Aria | 50mg/4ml | ☐ Initiation – Infuse 2mg/kg (mg) IV over 30 minutes at week 0, week 4, and week 8 | | | ☐ TB Skin Test ☐ CBC w/Diff q | | | | |
| | | ☐ Maintenance – Infuse 2mg/kg (mg) IV over 30 minutes every 8 weeks | | | | | | | |
| ☐ Simponi | □ 50mg/0.5ml Smartject | Inject 50mg SQ once monthly | | | ☐ TB Skin Test ☐ CBC w/Diff g | | | | |
| | ☐ 50mg/0.5ml PFS | | | | , , | | | | |
| ☐ Erivedge | 150mg Capsules | Take 1 (one) capsule by mouth daily | | | | | | | |
| ☐ Inflectra | | ☐ Initiation - Inject mg IV infusion at 0,2,6 weeks and then every 8 weeks thereafter (5mg/kg) | | | | | | | |
| | | ☐ Maintenance - Infuse mg IV every 8 weeks (5mg/kg) | | | | | | | |
| ☐ Renflexis | | ☐ Initiation - Inject mg IV infusion at 0,2,6 weeks and then every 8 weeks thereafter (5mg/kg) | | | | | | | |
| | | ☐ Maintenance - Infuse mg IV every 8 weeks (5 mg/kg) | | | | | | | |
| Premedication(s): | | | IV Access Flush Order: | | | | | | |
| □ Diphenhydramine 25-50 mg po- 25mg #2 per dose □ Acetaminophen 325-650 mg po- 325mg #2 per dose □ Methylprednisolone mg IV over mins □ Other: □ All infusion supplies necessary to administer the medication □ Anaphylaxis Kit | | | | | | | | | |
| By signing below, I certify that above therapy is medically necessary. Prescriber's Signature (SIGN BELOW) By signing this form and utilizing our services, I am also authorizing TwelveStone to serve as my prior authorization agent with medical and pharmacy insurance providers. | | | | | | | | | |
| | | | | | | | | | |
| Dispense as Written | | Printed Name | Substitution Allowed Date | te | | | | | |

The information contained in this facsimile may be confidential and is intended solely for the use of the named recipient(s). Access, copying or re-use of the facsimile or any information contained therein by any other person is not authorized. If you are not the intended recipient, please notify us immediately by calling 615-895-0186 or faxing back to the originator.