Blincyto Enrollment Form			TwelveStone Health Partners			
Date: Patient Name: Date of Birth:		D	Fax Refe (800) 22 irect Phone: (6 Toll Free: (844	3 -4063 515) 278-3350	TwelveStone HEALTH PARTNERS	
PREVIOUS ADMINISTRATION						
If YES, please provide the following information:				If NO, please indicate desired location for first dose:		
Last Infusion Date:				☐ Physician's Office		
Next Infusion Date:				☐ TwelveStone Infusion Center		
				☐ TwelveStone Home Infusion		
				☐ Other:		
				Desired Start Date:		
DIAGN						
Description ☐ B-Cell Precursor Acute Lymphoblastic Leukemia				ICD-10 Code □ C91.0		
	ОТІ	IER REQUIRED DOCU	MENTATION (PI	 lease attach documents	s as needed)	
☐ This signed order form ☐ History and Physical						
9 .					ab work (including most recent renal function tests porting primary diagnosis)	
CLINICAL INFORMATION (Please attach all clinical information, lab results and other medical history documents)						
Patient Weight:	:Kg/Lbs Hei	ght:Inches,	/CM BSA:	Allergies:		
Line Access: ☐ PIV ☐ PICC (SL DL TL) ☐ PORT (Huber sizeGaugeLength) ☐ Sub-Q						
MEDICATION	DOSE	DIRECTIONS			LAB & ANCILLARY ORDERS	
□ Blincyto	35mcg Vial: ☐ > 45kg (fixed dose) ☐ < 45kg (BSA based dose)	daily on days 10-28 interval on days 29- Cycle 2 - Infuse 28m daily on days 3-28; interval on days 29- E) Consolidation - Configure 28mcg (or 15 days 1-28; followed days 29-42 Continued Thera Infuse 28mcg (or 15 days 15 days 15 days 29-42	ncg (or 15mcg/m2; followed by 14 d 42 ncg (or 15mcg/m2 followed by 14 da 42 cycles 3-5 smcg/m2) continu by 14 days treatr py - Cycles 6-9 smcg/m2) continu	2) continuous IV infusion	□ Okay to proceed if ANC > and PLT > □ Adjust dose by % if ANC > and < □ Adjust dose by % if PLT > and < □ Hold dose if ANC < and/or PLT < *Will Notify MD about any dose reduction **If Dosing Parameters are not selected then MD will be contacted for any lab or result not in the normal range	
Premedication(s):				ncillary Orders:		
 □ Diphenhydramine 25-50 mg po – 25mg #2 per dose □ Acetaminophen 325-650 mg po – 325mg #2 per dose □ Methylprednisolonemg IV overmins □ Other: 				□ NaCl 0.9% 5-10ml IV before and after infusion □ Heparin 10 units/ml 3-5ml IV after infusion for peripheral access and PRN □ Heparin 100 units/ml 3-5ml IV after infusion for central IV access and PRN □ All infusion supplies necessary to administer the medication □ Anaphylaxis Kit		
	elow, I certify that ab form and utilizing our serv		-		ture (SIGN BELOW) rization agent with medical and pharmacy insurance	
Physician' Phone# Physicia		hysician's NPI#	ician's NPI# Phy		Physician's Address	
Dispense as Written		<u> </u>		ubstitution Allowed	Date	